


# Accepted Manuscript

Detection of the NS3 Q80K polymorphism by Sanger and deep sequencing in hepatitis C virus (HCV) genotype 1a strains in the United Kingdom

Apostolos Beloukas, Simon King, Kate Childs, Athanasios Papadimitropoulos, Mark Hopkins, Mark Atkins, Kosh Agarwal, Mark Nelson, Prof Anna Maria Geretti, MD, PhD



PII: S1198-743X(15)00734-X

DOI: [10.1016/j.cmi.2015.07.017](https://doi.org/10.1016/j.cmi.2015.07.017)

Reference: CMI 345

To appear in: *Clinical Microbiology and Infection*

Received Date: 9 April 2015

Revised Date: 14 July 2015

Accepted Date: 19 July 2015

Please cite this article as: Beloukas A, King S, Childs K, Papadimitropoulos A, Hopkins M, Atkins M, Agarwal K, Nelson M, Geretti AM, Detection of the NS3 Q80K polymorphism by Sanger and deep sequencing in hepatitis C virus (HCV) genotype 1a strains in the United Kingdom, *Clinical Microbiology and Infection* (2015), doi: 10.1016/j.cmi.2015.07.017.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Detection of the NS3 Q80K polymorphism by Sanger and deep sequencing in hepatitis C virus (HCV) genotype 1a strains in the United Kingdom**

Apostolos Beloukas<sup>1\*</sup>, Simon King<sup>1\*</sup>, Kate Childs<sup>2</sup>, **Athanasios** Papadimitropoulos<sup>1</sup>, Mark Hopkins<sup>3</sup>, Mark Atkins<sup>4</sup>, Kosh Agarwal<sup>5</sup>, Mark Nelson<sup>6</sup>, Anna Maria Geretti<sup>1</sup>

\*The two authors contributed equally to this work.

<sup>1</sup>Institute of Infection & Global Health, University of Liverpool, Liverpool; <sup>2</sup>Dept. of Sexual Health, King's College Hospital NHS Foundation Trust, London; <sup>3</sup>Liverpool Specialist Virology Centre, Royal Liverpool University Hospital, Liverpool; <sup>4</sup>Dept. of Microbiology, Frimley Park Hospital NHS Foundation Trust, Frimley, Surrey; <sup>5</sup>Institute of Liver Studies, King's College Hospital NHS Foundation Trust, London; <sup>6</sup>HIV and Sexual Health Services, Chelsea and Westminster Hospital, London, United Kingdom.

**Keywords:** NS3, Q80K, lineage, simeprevir, protease inhibitors

**Running title:** Prevalence of Q80K in HCV-1a carriers in the UK

**Word count:** 2500

**Abstract:** 250

**Corresponding author**

Prof Anna Maria Geretti, MD, PhD

Institute of Infection & Global Health, University of Liverpool,

8 West Derby Street, L69 7BE, United Kingdom

+44 151 795 9665; [geretti@liverpool.ac.uk](mailto:geretti@liverpool.ac.uk)

**ABSTRACT**

The Q80K polymorphism in the HCV NS3 enzyme reduces susceptibility to simeprevir and other novel protease inhibitors. The study aimed to determine the prevalence of Q80K in treatment-naïve HCV-1a carriers in the North-West (NW) and South-East (SE) of England, investigate occurrence of Q80K as a minority variant, and characterise viral phylogeny. Plasma samples from subjects that were naïve to anti-HCV therapy underwent conventional (Sanger) and deep (Illumina-Miseq, 1% interpretative cut-off) sequencing of NS3. Q80K occurred in 44/238 subjects (18.5%, 95% confidence interval 13.6%-23.4%), including 19/70 (27.1%) in the NW and 25/168 (14.9%) in the SE ( $p=0.0425$ ), with no difference by HCV RNA load or HIV status. **Q80K frequencies in reads of samples that underwent Illumina sequencing were >40% in all cases. Among subjects with Q80K, 5/44 (11.4%) showed one additional major resistance-associated mutation in NS3 detected at mutant frequency above (V36L, V55A) or below (V36M) 10%.** Phylogenetic analyses identified the two recognised HCV-1a lineages with (clade I) and without (clade II) Q80K. Overall, 148/238 (62.2%) sequences occurred within regional or inter-regional clusters, **each comprising 3-20 sequences.** There was no unique clustering of English sequences relative to strains from continental Europe and North America. In conclusion, Q80K was found at high prevalence among treatment-naïve HCV-1a carriers in England, and was reliably detected by conventional sequencing, **with no increased detection by deep sequencing.** English sequences were highly interspersed with sequences from elsewhere in Europe (clade II) and North America (clade I), and their phylogeny was consistent with multiple introductions from different areas.

## INTRODUCTION

The HCV NS3 protease enzyme exhibits a high degree of genetic variability and only 47% of its amino acids are conserved among circulating HCV genotypes.[1] NS3 genetic heterogeneity and the associated molecular and structural differences influence HCV susceptibility to protease inhibitors (PIs), including licensed first-generation (telaprevir, boceprevir) and second-generation (simeprevir) compounds. HCV variants with reduced PI susceptibility have been observed in treatment-naïve patients.[1-10] Among these, the glutamine to lysine substitution at codon 80 (Q80K) reduces susceptibility to simeprevir *in vitro* [2, 4, 11] and was also seen to reduce virological responses to simeprevir plus pegylated interferon alpha and ribavirin (P/R) in clinical studies.[12-14] Screening for Q80K is therefore recommended before starting simeprevir.[15]

The prevalence of Q80K among PI-naïve subjects varies by HCV genotype and subtype. An analysis of global NS3 sequences deposited in GenBank reported Q80K in 42% of HCV-1a strains, and also detected a high prevalence in HCV-5 and HCV-6 strains.[1] Among HCV-1a strains, prevalence of Q80K ranges between 3% and 47% by geographical location, exceeding 40% in North America.[2, 4] Reported prevalence is 20% overall across Europe, although differing markedly by country.[7-9] Most studies to date have produced prevalence estimates using conventional (Sanger) sequencing, which allows detection of virus variants present at a frequency  $\geq 20\%$  within a patient's sample.

This study had three aims. Firstly, to determine the prevalence of Q80K in NS3 sequences obtained from HCV-1a carriers attending for care in two regions of England in the United Kingdom(UK). A further aim was to investigate the occurrence of Q80K as a low frequency variant by deep sequencing, and thereby both gain insights into the viral quasispecies and determine whether conventional sequencing offers sufficient sensitivity for screening simeprevir candidates for the presence of Q80K. The third aim was to determine the phylogeny of NS3 sequences in relation to publically available sequences from the rest of Europe and North America.

## METHODS

### *Study population*

The study was performed retrospectively using stored plasma samples from consecutive adults that in 2006-2014 attended for care at the Royal Liverpool University Hospital in Liverpool (North-West region) and at King's College Hospital, Chelsea & Westminster Hospital, and Charing Cross Hospital in London (South-East region). Eligible patients were infected with HCV-1a **based on the genotyping method available at the local National Health System diagnostic laboratory; the assignment was confirmed by analysis of the NS3 sequences produced in this study.** Patients were naïve to all anti-HCV therapy. HIV status and HCV RNA load were retrieved from the clinics' databases. Ethics permission was granted by the South Berkshire Regional Ethics Committee to conduct the study after removing personal identifiers from the samples.

### *Sanger sequencing*

HCV RNA was extracted with Nuclisens easyMAG (bioMérieux, Netherlands) and cDNA synthesis was performed with the Qiagen OneStep RT-PCR Kit (Hilden, Germany) with forward primer 3278 [GGAGACCAAGMTCATCACSTGG] and reverse primer 4032 [GCTCTTRCCGCTGCCRGTTGGG]. The first-round amplicon was subjected to nested PCR using the Qiagen HotStarTaq Kit with forward primer 3307 [ACACCGCSGCGTGYGGKGACAT] and reverse primer 4014 [GGRGCRTGYAGRTGGGCCAC]. The second-round 727bp amplicon (aa180 of NS2 to aa204 of NS3) was purified with the Qiagen QIAquick PCR Purification Kit and sequenced using the BigDye Terminator Cycle Sequencing Kit v3.1 on the ABI Prism 3730 Genetic Analyser (Applied Biosystems®, USA). Consensus sequences were assembled using SeqScape (v2.7) and analysed with the geno2pheno system (<http://hcv.geno2pheno.org/index.php>; March 2015 version) for the presence of HCV resistance-associated mutations (RAMs).

### *Deep sequencing*

HCV RNA was extracted with QiasymphonySP (Qiagen) using the Qiagen DSPVirus/Pathogen Midi Kit and the CellFree\_500\_V4 protocol; cDNA was produced using the ImProm II Reverse Transcription System (Promega, USA) and random primers. NS3 was amplified using Q5 High-Fidelity DNA Polymerase (New England Biolabs, USA) and the above primers. PCR products were purified using AMPure XP Beads (Beckman Coulter, USA) before quantification with the Quant-iT dsDNA High Sensitivity Kit (Life Technologies, USA)

on a Qubit 2.0 Fluorometer (Life Technologies). The Nextera XT Sample Preparation Kit (Illumina, USA) was used for library preparation and pooled amplicon libraries were sequenced on Illumina MiSeq using v2 reagents. Data generation and initial analysis were carried out at the Centre for Genomic Research at the University of Liverpool. Adapter sequences were trimmed from the Illumina reads using Cutadapt v1.2.1[16]; further trimming was done using Sickle v1.2 (<http://github.com/najoshi/sickle>) applying a minimum window quality score of 20. BAMStats (<http://sourceforge.net/projects/bamstats/files/>) was used to calculate the read statistics. Variants were analysed using the VirVarSeq pipeline,[17] which utilises the quality of the run and individual bases to filter poor quality bases and reduce false positive rates. A 727bp region of a HCV-1a molecular clone encoding full-length NS3 was amplified and sequenced in four replicates in order to estimate the assay error rate. The plasmid control yielded an average of 174,799 reads, with a mean coverage of 46,268 reads per nucleotide. The error rate, calculated by counting **false substitution reads** at the codon level of the molecular clone [18], was mean 0.6% (SD 0.2%) over NS3 amino acids 1-181.

#### *Phylogenetic analysis*

Reference HCV-1a sequences and all available HCV-1a NS3 sequences from North America and Europe with confirmed genotype assignment and country of origin were retrieved from the Los Alamos HCV database ([hcv.lanl.gov/content/sequence/HCV/ToolsOutline.html](http://hcv.lanl.gov/content/sequence/HCV/ToolsOutline.html); accessed on 8 March 2015) (**supplementary file**). **All sequences, including those produced in this study (Sanger sequences and NGS consensus sequences with 1% interpretative cut-off),** were aligned pairwise against the HCV-1a reference genome H77 (accession no. NC\_004102) using MEGA v.6. The alignment **was trimmed** to codon positions 1-194 of NS3 (nucleotides 3421-4014 of H77). Sequences with gaps or ambiguous nucleotides affecting  $\geq 50\%$  of the targeted region or where codon 80 was missing or ambiguous were removed (**n=3**), yielding a total of 1162 NS3 sequences for analysis. Two maximum-likelihood (ML) phylogenies were estimated with and without the 80 codon, using a general-time reversible nucleotide substitution model with gamma-distributed among-site rate variation and the ML method as implemented in PHYML v.3.0.[19, 20] Phylogenetic support was evaluated using a bootstrap approach and a total of 1000 bootstrap replicates were generated using the FastTree v.2.1.7 software;[21] local branch support was calculated by the Shimodaira–Hasegawa-like (SH-like) test. Phylogenies were visualized and annotated in FigTree (<http://tree.bio.ed.ac.uk>). Clusters ( $\geq 3$  sequences) were identified by a bootstrap support  $>75\%$  and were defined as regional if

comprising sequences unique to either the NW or the SE, and inter-regional if including sequences from both regions.

### *Statistical analysis*

Descriptive statistics were used to analyse the prevalence of mutants in the study population (as the proportion of subjects showing the mutation) and the frequency of mutants in each patient's sample (as the proportion of deep sequencing reads showing the mutation). Proportion of subjects with Q80K by geographic location and HIV status were compared by Fisher's exact test. The Mann-Whitney U test was used to compare the HCV RNA load of patients with and without Q80K. The analysis was performed using SPSS v.21.

## **RESULTS**

### *Study population*

Overall 238 adults (median age 44 years, with no difference by region) infected with HCV-1a were studied, including 70/238 (29.4%) in the North-West (NW) and 168/238 (70.6%) in the South-East (SE). At the time of testing, HCV RNA load was median 6.3 log<sub>10</sub> IU/ml (interquartile range, IQR 5.8-6.8), without differences between the two regions. Subjects were naïve to all anti-HCV therapy. A total of 61/238 (25.6%) subjects, all from the SE, were co-infected with HIV.

### *Q80K prevalence by Sanger sequencing*

Overall, Q80K was detected in 44/238 subjects, yielding a prevalence of 18.5% (95% confidence interval, CI 13.6%-23.4%)(Table 1). Samples from two of the 44 subjects had a mixed first base at codon 80 (Q80Q/K). **The median HCV RNA load was 6.4 (IQR 5.8-6.7) vs. 6.3 (IQR 5.8-6.8) log<sub>10</sub> IU/ml in samples with vs. samples without Q80K, respectively (p=0.63).** The prevalence of Q80K was 19/70 (27.1%) in the NW and 25/168 (14.9%) in the SE (p=0.043). When comparing results by HIV status in the SE, Q80K prevalence was 18/107 (16.8%) in HCV mono-infected subjects and 7/61 (11.5%) in HCV/HIV co-infected subjects (p=0.379). Table 1 shows other major and minor NS3 RAMs detected by Sanger sequencing.

### *Q80K prevalence by deep sequencing*

A total of 178/238 (74.8%) samples underwent deep sequencing, comprising 28/178 (15.7%) with and 150/178 (84.3%) without Q80K by Sanger sequencing. An average of 58,585 reads per sample was obtained (median 53,413; IQR 40,740-70,255) with an average coverage per



nucleotide of 16,107. Prevalence of Q80K was 27/178 (15.2% 95% CI 9.9%-20.5%). All 28 samples showing Q80K by Sanger sequencing also showed the mutation by deep sequencing, usually (26/28 samples) with mutant frequencies of  $\geq 98\%$ . Of the two samples showing Q80Q/K by Sanger sequencing, one had a Q:K ratio of 54:46, and the other showed Q, K and L at a ratio of 57:41:2. Of 150 samples lacking Q80K by Sanger sequencing, none showed the mutation by deep sequencing when applying an interpretative cut-off of  $\geq 1\%$  for the frequency of mutants within the reads, as usually recommended.[22] Prevalence of Q80K increased when the interpretative cut-off was lowered, and was 3/150 (2%) for Q80K occurring at a frequency between  $\geq 0.5\%$  and  $< 1\%$ , and 15/150 (10%) at a frequency  $\geq 0.2\%$  and  $< 0.5\%$ . These cut-offs however fell within the estimated error rate of the assay (0.6%, SD 0.2%).

Deep sequencing (interpretative cut-off  $\geq 1\%$ ) identified additional NS3 RAMs that had not been observed by Sanger sequencing and which occurred at frequencies between 1% and 6% (Table 1). **Among the 44 subjects with Q80K, 4/44 (9.1%) had one additional major NS3 RAM at frequency  $> 10\%$  including V36L in 3/44 (6.8%) and V55A in 1/44 (2.3%). In addition 1/44 (2.3%) subjects had V36M detected only by Illumina at a frequency of 3%.**

#### *Phylogeny of NS3 sequences*

HCV-1a strains separated into the two recognised distinct lineages with and without Q80K.[23, 24] The sequences harbouring Q80K at frequency  $< 1\%$  did not align with the Q80K lineage (data not shown). There was no unique clustering of the study English sequences (Figure 1), rather the sequences were interspersed with sequences from the rest of Europe and North America, and the phylogeny was consistent with multiple introductions from different areas. With the English sequences, four NW, eight SE, and 11 inter-regional clusters were identified, each consisting of 3-20 sequences (Figure 2). In total, 148/238 (62.2%) sequences (49/70, 70.0% in the NW and 99/168, 58.9% in the SE) were circulating as regional or inter-regional clusters.

## **DISCUSSION**

This study detected the NS3 polymorphism Q80K at high prevalence among treatment-naïve HCV-1a carriers attending for care in England, particularly in the North-West where prevalence reached 27.1%. Previously published data from the UK include a small cohort of 38 HIV-positive patients with acute HCV-1a infection, of which 16% had Q80K.[6] A larger study analysed 159 subjects with HCV-1a that were enrolled in Phase II/III studies of



telaprevir and simeprevir, of which 23% showed Q80K[8]; these sequences are not available for analysis however, as to date they have not been deposited in the GenBank or the Los Alamos database. These data indicate that HCV-1a strains circulating in the UK have the highest prevalence of Q80K observed in Europe.

In this study the prevalence of Q80K varied by geographical region, while showing no difference by HIV status. Prevalence rates also show marked geographical variability in the rest of Europe.[7-9] The variable prevalence has not been linked to race or ethnicity [8], but rather is consistent with the circulation of two distinct HCV-1a lineages with and without Q80K.[23, 24] The Q80K-carrying lineage is believed to have originated in the United States in the 1940s.[23] In this study, the maximum likelihood phylogenetic analysis confirmed the two HCV-1a lineages, and detected multiple introductions of the two lineages in the UK, with wide interspersing of the English strains with other European and North American strains. Most HCV-1a sequences occurred as regional or mixed regional clusters, however, suggesting that for this cohort the majority of transmission events occurred within the UK.

The deep sequencing analysis showed that Q80K mutants occurred at high frequency (>40%) in the patients' samples, allowing detection by conventional sequencing in all cases. This finding is consistent with the observation that Q80K mutants have replication capacity similar to that of wild-type virus, which allows transmissibility and persistence at high frequency within the viral quasispecies despite the absence of drug selective pressure. There were no samples showing Q80K at a frequency below the detection threshold of Sanger sequencing ( $\geq 10\text{-}20\%$ ) and above the typical  $\geq 1\%$  interpretative cut-off for deep sequencing, which is recommended to differentiate biologically meaningful mutations from those caused by methodological errors.[22] When applying less stringent cut-offs (0.5% or 0.2%) the proportion of samples showing Q80K by deep sequencing increased. A previous study similarly found that among 21 subjects with Q80K detected by deep sequencing, four showed the mutation at a frequency  $< 1\%$ .[10] The estimated error rate of the assay (0.6%) and the observation that sequences showing Q80K at a frequency  $< 1\%$  did not align within the Q80K lineage suggest that the detection of Q80K at frequency  $< 1\%$  was spurious. **Furthermore, it remains to be determined whether any Q80K variant occurring at low frequency has clinical relevance.**

Q80K reduces HCV-1a susceptibility to simeprevir by 8-11 fold *in vitro*. [4, 8, 11] In HCV-1a infection, sustained virological responses (SVR) to combination therapy with simeprevir plus P/R are reduced by the presence of baseline Q80K in both treatment-naïve subjects (from 84% to 58%) and prior relapsers (from 79% to 47%). [11-13] The impact of Q80K on the combination of simeprevir plus sofosbuvir was not immediately apparent in some reported studies. [25] **However, a phase III, open-label, single-arm study with simeprevir plus sofosbuvir in cirrhotic patients with HCV-1 recently reported that SVR rates were higher in subjects without Q80K (92%) versus those with the mutation (74%).** [26] Q80K has no effects on susceptibility to telaprevir and boceprevir [4], but may have an effect on responses to asunaprevir. [27] Q80K also confers small reductions (~3-fold) in susceptibility to faldaprevir and paritaprevir *in vitro*, without necessarily affecting response rates. [28] The Q80L and Q80R mutations were detected in some subjects, usually at low frequency. Their significance is unclear. *In vitro*, the mutations confer 2-fold and 7-fold reductions in susceptibility to simeprevir, respectively. [11] Q80R has also been observed in a small number of patients experiencing failure of simeprevir therapy, when it occurred alongside major resistance mutations at position 155 and/or position 168. [29] Other major NS3 RAMs occurred at codons 36, 54, 55, 168, and 170, which is consistent with previous observations. [1-6, 9, 30]. **A small subset of five subjects had Q80K plus one other major RAM in NS3.**

In summary, we found a high prevalence of Q80K in HCV-1a carriers with and without HIV accessing care in two regions of the UK, thus complementing estimates from elsewhere in Europe. While deep sequencing increased detection of Q80K, mutants missed by Sanger sequencing occurred at frequency  $\leq 0.5\%$  and detection seemed a likely technical artefact. There was evidence of a high degree of interspersing of UK sequences with sequences from elsewhere in Europe and North America, although the phylogeny indicated that most transmission events occurred in the UK. Investigation of full viral genomes will elucidate HCV-1a transmission networks and gain data to inform control strategies.


**Table 1.** Prevalence of NS3 resistance-associated mutations (RAMs) in treatment-naïve HCV-1a carriers according to modality of detection (Sanger or Illumina) and frequency of the mutant as determined by Illumina<sup>a</sup>

RAMs	Sanger/Illumina >10% n (%)	Illumina 1-10% n (%)
Total tested	238 (100)	178 (100)
<b>36 M</b>	1 (0.4)	2 (1.1)
<b>36 L</b>	5 (2.1)	0 (0)
<b>54 S</b>	9 (3.8)	0 (0)
<b>55 A</b>	10 (4.2)	0 (0)
<b>80 K</b>	44 (18.5)	0 (0)
80 L	1 (0.4)	4 (2.2)
80 R	0 (0)	3 (1.7)
117 H	0 (0)	1 (0.6)
<b>168 E</b>	1 (0.4)	0 (0)
<b>170 A</b>	1 (0.4)	0 (0)
<b>170 T</b>	0 (0)	2 (1.1)
174 S	97 (40.8)	10 (5.6)

<sup>a</sup>RAMs were classified according to the geno2pheno mutation list (March 2015 version) with the addition of Q80L (reference 11). Bold indicates major mutations for genotype 1/1a.

**Figure 1.** Phylogenetic analysis of HCV-1a NS3 sequences from the United Kingdom, where green indicates sequences from the North-West and yellow indicates sequences from the South-East of England. The global reference dataset was derived from Los Alamos HCV database (March 2015; see supplementary file for accession numbers) and comprises sequences from the rest of Europe (in red) and North America (in blue). Nodes with bootstrap support values  $\geq 90\%$  are indicated with an asterisk.


**80 K phylogeny**



**80 Q phylogeny**

284 **Figure 2.** A HCV transmission cluster in the South-East of England, comprising 20 HCV-1a  
 285 NS3 sequences (in yellow). Sequences from the rest of Europe and North America are shown  
 286 in red and blue respectively. Nodes with bootstrap support values  $\geq 90\%$  are indicated with an  
 287 asterisk.

288



ACCEPTED

## Acknowledgements

The work was supported by research awards from the British HIV Association and the Rosetrees Trust in the United Kingdom.

**Part of this work was presented at the 13<sup>th</sup> European Meeting on HIV & Hepatitis (Barcelona, Spain; June 2015) and at the 11<sup>th</sup> International Workshop on HIV & Hepatitis Co-infection (London, UK; June 2015)**

## References

- 1 Cento V, Mirabelli C, Salpini R, et al. Hcv genotypes are differently prone to the development of resistance to linear and macrocyclic protease inhibitors. *PloS one*. 2012; **7**: e39652.
- 2 Bae A, Sun SC, Qi X, et al. Susceptibility of treatment-naïve hepatitis c virus (HCV) clinical isolates to hcv protease inhibitors. *Antimicrobial Agents and Chemotherapy*. 2010; **54**: 5288-5297.
- 3 Bartels DJ, Sullivan JC, Zhang EZ, et al. Hepatitis c virus variants with decreased sensitivity to direct-acting antivirals (daas) were rarely observed in daa-naïve patients prior to treatment. *Journal of Virology*. 2013; **87**: 1544-1553.
- 4 Berger KL, Triki I, Cartier M, et al. Baseline hepatitis c virus (hcv) ns3 polymorphisms and their impact on treatment response in clinical studies of the hcv ns3 protease inhibitor faldaprevir. *Antimicrobial Agents and Chemotherapy*. 2014; **58**: 698-705.
- 5 Kuntzen T, Timm J, Berical A, et al. Naturally occurring dominant resistance mutations to hepatitis c virus protease and polymerase inhibitors in treatment-naïve patients. *Hepatology*. 2008; **48**: 1769-1778.
- 6 Leggewie M, Sreenu VB, Abdelrahman T, et al. Natural ns3 resistance polymorphisms occur frequently prior to treatment in hiv-positive patients with acute hepatitis c. *AIDS*. 2013; **27**: 2485-2488.
- 7 Morel V, Duverlie G, Brochot E. Patients eligible for treatment with simeprevir in a french center. *Journal of Clinical Virology*. 2014; **61**: 149-151.
- 8 Sarrazin C, Lathouwers E, Peeters M, et al. Prevalence of the hepatitis c virus ns3 polymorphism q80k in genotype 1 patients in the european region. *Antiviral Research*. 2015; **116C**: 10-16.
- 9 Vicenti I, Rosi A, Saladini F, et al. Naturally occurring hepatitis c virus (HCV) ns3/4a protease inhibitor resistance-related mutations in HCV genotype 1-infected subjects in italy. *Journal of Antimicrobial Chemotherapy*. 2012; **67**: 984-987.
- 10 Jabara CB, Hu F, Mollan KR, et al. Hepatitis c virus (HCV) NS3 sequence diversity and antiviral resistance-associated variant frequency in HCV/HIV coinfection. *Antimicrobial Agents and Chemotherapy*. 2014; **58**: 6079-6092.
- 11 Lenz O, Verbinen T, Lin TI, et al. In vitro resistance profile of the hepatitis c virus ns3/4a protease inhibitor tmc435. *Antimicrobial Agents and Chemotherapy*. 2010; **54**: 1878-1887.
- 12 Forns X, Lawitz E, Zeuzem S, et al. Simeprevir with peginterferon and ribavirin leads to high rates of svr in patients with hcv genotype 1 who relapsed after previous therapy: A phase 3 trial. *Gastroenterology*. 2014; **146**: 1669-1679 e1663.
- 13 Jacobson IM, Dore GJ, Foster GR, et al. Simeprevir with pegylated interferon alfa 2a plus ribavirin in treatment-naïve patients with chronic hepatitis c virus genotype 1



- infection (quest-1): A phase 3, randomised, double-blind, placebo-controlled trial. *Lancet*. 2014; **384**: 403-413.
- 14 Manns M, Marcellin P, Poordad F, et al. Simeprevir with pegylated interferon alfa 2a or 2b plus ribavirin in treatment-naïve patients with chronic hepatitis c virus genotype 1 infection (quest-2): A randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2014; **384**: 414-426.
- 15 Simeprevir summary of product characteristics. Available at: [Http://www.Ema.Europa.Eu/docs/en\\_gb/document\\_library/epar\\_-\\_product\\_information/human/002777/wc500167867.Pdf](http://www.Ema.Europa.Eu/docs/en_gb/document_library/epar_-_product_information/human/002777/wc500167867.Pdf). (accessed 3 march 2015).
- 16 M. M. Cutadapt removes adapter sequences from high-throughput sequencing reads. . *EMBnet J* 2011; **17**: 10-12.
- 17 Verbist BM, Thys K, Reumers J, et al. Virvarseq: A low-frequency virus variant detection pipeline for illumina sequencing using adaptive base-calling accuracy filtering. *Bioinformatics*. 2015; **31**: 94-101.
- 18 Dierynck I, Thys K, Ghys A, et al. Deep-sequencing analysis of the gene encoding the hepatitis c virus nonstructural 3-4a protease confirms a low prevalence of telaprevir-resistant variants at baseline and the end of the realize study. *Journal of Infectious Diseases*. 2014; **210**: 1871-1880.
- 19 Beloukas A, Magiorkinis E, Magiorkinis G, et al. Assessment of phylogenetic sensitivity for reconstructing hiv-1 epidemiological relationships. *Virus Res*. 2012; **166**: 54-60.
- 20 Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of phyml 3.0. *Systematic Biology*. 2010; **59**: 307-321.
- 21 Price MN, Dehal PS, Arkin AP. Fasttree 2--approximately maximum-likelihood trees for large alignments. *PloS One*. 2010; **5**: e9490.
- 22 Gianella S, Delport W, Pacold ME, et al. Detection of minority resistance during early hiv-1 infection: Natural variation and spurious detection rather than transmission and evolution of multiple viral variants. *Journal of Virology*. 2011; **85**: 8359-8367.
- 23 McCloskey RM, Liang RH, Joy JB, et al. Global origin and transmission of hepatitis c virus nonstructural protein 3 q80k polymorphism. *Journal of Infectious Diseases*. 2014.
- 24 Pickett BE, Striker R, Lefkowitz EJ. Evidence for separation of hcv subtype 1a into two distinct clades. *Journal of Viral Hepatitis*. 2011; **18**: 608-618.
- 25 **Lawitz E, Sulkowski MS, Ghalib R, et al. Simeprevir plus sofosbuvir, with or without ribavirin, to treat chronic infection with hepatitis c virus genotype 1 in non-responders to pegylated interferon and ribavirin and treatment-naïve patients: The cosmos randomised study. *Lancet*. 2014; **384**: 1756-1765.**
- 26 **Lawitz E, Matusow G, DeJesus E. A phase 3, open-label, single-arm study to evaluate the efficacy and safety of 12 weeks of simeprevir (smv) plus sofosbuvir (sof) in treatment-naïve or experienced patients with chronic HCV genotype 1 (HCV-1) infection and cirrhosis: Optimist-2. 50th Annual Meeting of the European Association for the Study of the Liver. Vienna, Austria - April 22-26, 2015.**
- 27 McPhee F, Friborg J, Levine S, et al. Resistance analysis of the hepatitis c virus NS3 protease inhibitor asunaprevir. *Antimicrobial Agents and Chemotherapy*. 2012; **56**: 3670-3681.
- 28 Pilot-Matias T, Tripathi R, Cohen D, et al. In vitro and in vivo antiviral activity and resistance profile of the hepatitis c virus NS3/4a protease inhibitor abt-450. *Antimicrobial Agents and Chemotherapy*. 2015; **59**: 988-997.



- 384 29 Lenz O, Verbinnen T, Fevery B, et al. Virology analyses of HCV isolates from  
385 genotype 1-infected patients treated with simeprevir plus peginterferon/ribavirin in  
386 phase iib/iii studies. *Journal of Hepatology*. 2014.
- 387 30 Akuta N, Suzuki F, Sezaki H, et al. Evolution of simeprevir-resistant variants over time  
388 by ultra-deep sequencing in hcv genotype 1b. *Journal of Medical Virology*. 2014; **86**:  
389 1314-1322.

390

AF009606

AY695436

AF011751

AF011752

AF011753

AY615798

AY695437

AY956463

AY956464

AY956465

AY956466

AY956468

AY956469

DQ068178

DQ068179

DQ068180

DQ068181

DQ068182

DQ068183

DQ068184

DQ068185

DQ068186

DQ068187

DQ068188

DQ068189

DQ068190

DQ068191

DQ068192

DQ068193

DQ068194

DQ068195

DQ068196

DQ068197

DQ068198

DQ068199

DQ068200

DQ068201

DQ068202

DQ068203

DQ068204

DQ068205

DQ068206

DQ068207

DQ430811

DQ430812

DQ430813

DQ430814

DQ838739

DQ838740

DQ838741

DQ838742  
DQ838743  
DQ838744  
DQ838745  
DQ889251  
DQ889252  
DQ889253  
DQ889254  
DQ889255  
DQ889256  
DQ889257  
DQ889258  
DQ889259  
DQ889260  
DQ889261  
DQ889262  
DQ889263  
DQ889264  
DQ889265  
DQ889266  
DQ889267  
DQ889268  
DQ889269  
DQ889270  
DQ889271  
DQ889272  
DQ889273  
DQ889274  
DQ889275  
DQ889276  
DQ889277  
DQ889278  
DQ889279  
DQ889280  
DQ889281  
DQ889282  
DQ889283  
DQ889284  
DQ889286  
DQ889287  
DQ889288  
DQ889289  
DQ889290  
DQ889291  
DQ889292  
DQ889293  
DQ889294  
DQ889295  
DQ889296  
DQ889297

DQ889298

DQ889299

DQ889300

DQ889301

DQ889302

DQ889303

DQ889304

DQ889305

DQ889306

DQ889307

DQ889308

DQ889309

DQ889310

DQ889311

DQ889312

DQ889313

DQ889314

DQ889315

DQ889316

DQ889317

DQ889318

DQ889319

DQ889320

DQ889321

EF032883

EF032884

EF032885

EF139707

EF139710

EF139715

EF139719

EF139723

EF139725

EF139728

EF139729

EF139730

EF139735

EF139736

EF139740

EF139741

EF139746

EF139748

EF139749

EF139754

EF173627

EF173628

EF173629

EF173630

EF173631

EF173632

EF173633  
EF173634  
EF173635  
EF173636  
EF173637  
EF173638  
EF173639  
EF173640  
EF173641  
EF173642  
EF173643  
EF173644  
EF173645  
EF173646  
EF173647  
EF173648  
EF173649  
EF173650  
EF173651  
EF173652  
EF173653  
EF173654  
EF407411  
EF407412  
EF407413  
EF407414  
EF407415  
EF407416  
EF407417  
EF407418  
EF407419  
EF407420  
EF407421  
EF407422  
EF407423  
EF407424  
EF407425  
EF407426  
EF407427  
EF407428  
EF407429  
EF407430  
EF407431  
EF407432  
EF407433  
EF407434  
EF407435  
EF407436  
EF407437  
EF407438

EF407439  
EF407440  
EF407441  
EF407442  
EF407443  
EF407444  
EF407445  
EF407446  
EF407447  
EF407448  
EF407449  
EF407450  
EF407451  
EF407452  
EF407453  
EF407454  
EF407455  
EF407456  
EF407457  
EF621489  
EU155213  
EU155214  
EU155215  
EU155216  
EU155233  
EU155236  
EU155237  
EU155238  
EU155239  
EU155240  
EU155241  
EU155242  
EU155243  
EU155244  
EU155245  
EU155246  
EU155247  
EU155248  
EU155249  
EU155250  
EU155251  
EU155252  
EU155265  
EU155266  
EU155267  
EU155268  
EU155269  
EU155270  
EU155271  
EU155272

EU155273  
EU155274  
EU155275  
EU155276  
EU155277  
EU155278  
EU155282  
EU155283  
EU155284  
EU155285  
EU155286  
EU155287  
EU155288  
EU155289  
EU155290  
EU155291  
EU155292  
EU155293  
EU155294  
EU155295  
EU155296  
EU155297  
EU155298  
EU155299  
EU155309  
EU155310  
EU155311  
EU155312  
EU155313  
EU155314  
EU155319  
EU155320  
EU155321  
EU155322  
EU155323  
EU155338  
EU155339  
EU155340  
EU155341  
EU155342  
EU155343  
EU155344  
EU155345  
EU155346  
EU155347  
EU155348  
EU155349  
EU155350  
EU155351  
EU155352



EU155353  
EU155354  
EU155355  
EU155378  
EU155379  
EU155380  
EU234063  
EU234064  
EU234065  
EU239713  
EU239715  
EU239716  
EU250017  
EU255927  
EU255928  
EU255929  
EU255930  
EU255931  
EU255932  
EU255933  
EU255934  
EU255935  
EU255936  
EU255937  
EU255938  
EU255939  
EU255940  
EU255941  
EU255942  
EU255943  
EU255944  
EU255945  
EU255946  
EU255947  
EU255948  
EU255949  
EU255950  
EU255951  
EU255952  
EU255953  
EU255954  
EU255955  
EU255956  
EU255957  
EU255958  
EU255959  
EU255963  
EU255964  
EU255965  
EU255966

EU255967  
EU255968  
EU255969  
EU255970  
EU255971  
EU255973  
EU255974  
EU255975  
EU255976  
EU255977  
EU255978  
EU255979  
EU255980  
EU255981  
EU255982  
EU255983  
EU255984  
EU255985  
EU255986  
EU255987  
EU255988  
EU255989  
EU255990  
EU255991  
EU255992  
EU255993  
EU255994  
EU255995  
EU255996  
EU255997  
EU255998  
EU255999  
EU256002  
EU256003  
EU256004  
EU256005  
EU256006  
EU256007  
EU256008  
EU256009  
EU256010  
EU256011  
EU256012  
EU256013  
EU256014  
EU256015  
EU256016  
EU256017  
EU256018  
EU256019

EU256020  
EU256021  
EU256022  
EU256023  
EU256024  
EU256025  
EU256026  
EU256027  
EU256028  
EU256029  
EU256030  
EU256031  
EU256032  
EU256033  
EU256034  
EU256035  
EU256036  
EU256037  
EU256038  
EU256039  
EU256040  
EU256041  
EU256042  
EU256043  
EU256044  
EU256046  
EU256047  
EU256048  
EU256049  
EU256050  
EU256051  
EU256052  
EU256053  
EU256055  
EU256056  
EU256057  
EU256058  
EU256060  
EU256067  
EU256068  
EU256069  
EU256070  
EU256071  
EU256072  
EU256073  
EU256074  
EU256086  
EU256087  
EU256094  
EU256095

EU256096  
EU256097  
EU256104  
EU256105  
EU256106  
EU256107  
EU260395  
EU260396  
EU362876  
EU362877  
EU362878  
EU362879  
EU362880  
EU362881  
EU362882  
EU362883  
EU362884  
EU362885  
EU362886  
EU362887  
EU362888  
EU362889  
EU362890  
EU362891  
EU362892  
EU362893  
EU362894  
EU362895  
EU362896  
EU362897  
EU362898  
EU362899  
EU362900  
EU362901  
EU362902  
EU362903  
EU362904  
EU362905  
EU362906  
EU362907  
EU362908  
EU362909  
EU362910  
EU362911  
EU482831  
EU482832  
EU482834  
EU482835  
EU482836  
EU482837

EU482838  
EU482840  
EU482841  
EU482842  
EU482843  
EU482845  
EU482846  
EU482847  
EU482848  
EU482850  
EU482852  
EU482853  
EU482854  
EU482855  
EU482856  
EU482857  
EU482858  
EU482861  
EU482862  
EU482863  
EU482864  
EU482865  
EU482866  
EU482867  
EU482868  
EU482869  
EU482870  
EU482871  
EU482872  
EU482873  
EU482876  
EU482878  
EU482882  
EU482884  
EU482887  
EU482889  
EU529676  
EU529677  
EU529678  
EU529679  
EU529680  
EU529681  
EU569722  
EU569723  
EU595697  
EU595698  
EU595699  
EU622930  
EU660383  
EU660384

EU660385  
EU660387  
EU677247  
EU677248  
EU677249  
EU677250  
EU677251  
EU677252  
EU677253  
EU677254  
EU677255  
EU677256  
EU677257  
EU677258  
EU687193  
EU687194  
EU687195  
EU781746  
EU781747  
EU781748  
EU781749  
EU781750  
EU781751  
EU781752  
EU781753  
EU781754  
EU781755  
EU781756  
EU781757  
EU781758  
EU781759  
EU781760  
EU781761  
EU781762  
EU781763  
EU781764  
EU781765  
EU781766  
EU781767  
EU781768  
EU781769  
EU781770  
EU781771  
EU781772  
EU781773  
EU781774  
EU781775  
EU781776  
EU781777  
EU781778

EU781779  
EU781781  
EU781782  
EU781783  
EU781784  
EU781785  
EU781786  
EU781787  
EU781788  
EU781789  
EU781790  
EU781791  
EU781792  
EU781793  
EU781794  
EU781795  
EU781796  
EU781797  
EU781798  
EU781799  
EU781800  
EU781801  
EU781802  
EU781803  
EU781804  
EU862823  
EU862824  
EU862826  
EU862827  
EU862828  
EU862829  
EU862830  
EU862831  
EU862832  
EU862833  
EU862834  
EU862836  
EU862838  
EU862839  
EU862840  
EU862841  
FJ024087  
FJ024274  
FJ024275  
FJ024276  
FJ024278  
FJ024280  
FJ024281  
FJ024282  
FJ181999



FJ182000  
FJ182001  
FJ205867  
FJ205868  
FJ205869  
FJ390394  
FJ390395  
FJ390399  
FJ410172  
GQ149768  
HM568417  
HM568418  
HM568419  
HM568420  
HM568421  
HM568422  
HM568423  
HM568424  
HM568425  
HM746870  
HM746873  
HM746876  
HM746877  
HM746878  
HM746879  
HM746880  
HM746882  
JN704193  
JN704194  
JN704198  
JN704199  
JN704202  
JN704203  
JN704204  
JN704206  
JN704208  
JN704210  
JN704218  
JN704219  
JN704220  
JN704224  
JN704229  
JN704230  
JN704231  
JN704232  
JN704233  
JN704234  
JN704235  
JN704236  
JN704237

JN704238  
JN704239  
JN704240  
JN704241  
JN704242  
JN704243  
JN704244  
JN704245  
JN704246  
JN704247  
JN704248  
JN704249  
JN704250  
JN704251  
JN704252  
JN704253  
JN704254  
JN704255  
JN704256  
JN704257  
JN704260  
JN704261  
JN704262  
JN704263  
JN704264  
JN704266  
JN704267  
JN704268  
JN704269  
JN704270  
JN704273  
JN704274  
JN704275  
JN704276  
JN704277  
JN704278  
JN704280  
JN704281  
JN704282  
JN704283  
JN704284  
JN704285  
JN704290  
JN704291  
JN704293  
JN704294  
JN704295  
JQ061779  
JQ061780  
JQ061781

JQ061782  
JQ061783  
JQ061784  
JQ061785  
JQ061786  
JQ061787  
JQ914271  
JQ914273  
JQ914274  
JX463525  
JX463526  
JX463527  
JX463528  
JX463529  
JX463530  
JX463531  
JX463532  
JX463533  
JX463534  
JX463535  
JX463536  
JX463537  
JX463538  
JX463539  
JX463540  
JX463541  
JX463542  
JX463543  
JX463544  
JX463545  
JX463546  
JX463547  
JX463548  
JX463549  
JX463550  
JX463551  
JX463552  
JX463553  
JX463554  
JX463555  
JX463556  
JX463557  
JX463558  
JX463559  
JX463560  
JX463561  
JX463562  
JX463563  
JX463564  
JX463565

JX463566  
JX463567  
JX463568  
JX463569  
JX463570  
JX463571  
JX463572  
JX463573  
JX463574  
JX463575  
JX463576  
JX463577  
JX463578  
JX463579  
JX463580  
JX463581  
JX463582  
JX463583  
JX463584  
JX463585  
JX463586  
JX463587  
JX463588  
JX463589  
JX463590  
JX463591  
JX463592  
JX463593  
JX463594  
JX463595  
JX463596  
JX463597  
JX463598  
JX463599  
JX463600  
JX463601  
JX463602  
JX463603  
JX463604  
JX463605  
JX463606  
JX463607  
JX463608  
JX463609  
JX463610  
JX463611  
JX463612  
JX463613  
JX463614  
JX463615

JX463616  
JX463617  
JX463618  
JX463619  
JX463620  
JX463621  
JX463622  
JX463623  
JX463624  
JX463625  
JX463626  
JX463627  
JX463628  
JX463629  
JX463630  
JX463631  
JX463632  
JX463633  
JX463634  
JX463635  
JX463636  
JX463637  
JX463638  
JX463639  
JX463640  
JX463641  
KC155254  
M62321  
M67463  
NC\_004102